

Dietary Reference Intakes (DRIs): Recommended Dietary Allowances and Adequate Intakes, Vitamins

Food and Nutrition Board, Institute of Medicine, National Academies

Life Stage Group	Vitamin A (µg/d) ^a	Vitamin C (mg/d)	Vitamin D (µg/d) ^{b,c}	Vitamin E (mg/d) ^d	Vitamin K (µg/d)	Thiamin (mg/d)	Riboflavin (mg/d)	Niacin (mg/d) ^e	Vitamin B ₆ (mg/d)	Folate (µg/d) ^f	Vitamin B ₁₂ (µg/d)	Pantothenic Acid (mg/d)	Biotin (µg/d)	Choline (mg/d) ^g
Infants														
0 to 6 mo	400*	40*	10	4*	2.0*	0.2*	0.3*	2*	0.1*	65*	0.4*	1.7*	5*	125*
6 to 12 mo	500*	50*	10	5*	2.5*	0.3*	0.4*	4*	0.3*	80*	0.5*	1.8*	6*	150*
Children														
1–3 y	300	15	15	6	30*	0.5	0.5	6	0.5	150	0.9	2*	8*	200*
4–8 y	400	25	15	7	55*	0.6	0.6	8	0.6	200	1.2	3*	12*	250*
Males														
9–13 y	600	45	15	11	60*	0.9	0.9	12	1.0	300	1.8	4*	20*	375*
14–18 y	900	75	15	15	75*	1.2	1.3	16	1.3	400	2.4	5*	25*	550*
19–30 y	900	90	15	15	120*	1.2	1.3	16	1.3	400	2.4	5*	30*	550*
31–50 y	900	90	15	15	120*	1.2	1.3	16	1.3	400	2.4	5*	30*	550*
51–70 y	900	90	15	15	120*	1.2	1.3	16	1.7	400	2.4^h	5*	30*	550*
> 70 y	900	90	20	15	120*	1.2	1.3	16	1.7	400	2.4^h	5*	30*	550*
Females														
9–13 y	600	45	15	11	60*	0.9	0.9	12	1.0	300	1.8	4*	20*	375*
14–18 y	700	65	15	15	75*	1.0	1.0	14	1.2	400ⁱ	2.4	5*	25*	400*
19–30 y	700	75	15	15	90*	1.1	1.1	14	1.3	400ⁱ	2.4	5*	30*	425*
31–50 y	700	75	15	15	90*	1.1	1.1	14	1.3	400ⁱ	2.4	5*	30*	425*
51–70 y	700	75	15	15	90*	1.1	1.1	14	1.5	400	2.4^h	5*	30*	425*
> 70 y	700	75	20	15	90*	1.1	1.1	14	1.5	400	2.4^h	5*	30*	425*
Pregnancy														
14–18 y	750	80	15	15	75*	1.4	1.4	18	1.9	600ⁱ	2.6	6*	30*	450*
19–30 y	770	85	15	15	90*	1.4	1.4	18	1.9	600ⁱ	2.6	6*	30*	450*
31–50 y	770	85	15	15	90*	1.4	1.4	18	1.9	600ⁱ	2.6	6*	30*	450*
Lactation														
14–18 y	1,200	115	15	19	75*	1.4	1.6	17	2.0	500	2.8	7*	35*	550*
19–30 y	1,300	120	15	19	90*	1.4	1.6	17	2.0	500	2.8	7*	35*	550*
31–50 y	1,300	120	15	19	90*	1.4	1.6	17	2.0	500	2.8	7*	35*	550*

NOTE: This table (taken from the DRI reports, see www.nap.edu) presents Recommended Dietary Allowances (RDAs) in **bold type** and Adequate Intakes (AIs) in ordinary type followed by an asterisk (*). An RDA is the average daily dietary intake level; sufficient to meet the nutrient requirements of nearly all (97-98 percent) healthy individuals in a group. It is calculated from an Estimated Average Requirement (EAR). If sufficient scientific evidence is not available to establish an EAR, and thus calculate an RDA, an AI is usually developed. For healthy breastfed infants, an AI is the mean intake. The AI for other life stage and gender groups is believed to cover the needs of all healthy individuals in the groups, but lack of data or uncertainty in the data prevent being able to specify with confidence the percentage of individuals covered by this intake.

^a As retinol activity equivalents (RAEs). 1 RAE = 1 µg retinol, 12 µg β-carotene, 24 µg α-carotene, or 24 µg β-cryptoxanthin. The RAE for dietary provitamin A carotenoids is two-fold greater than retinol equivalents (RE), whereas the RAE for preformed vitamin A is the same as RE.

^b As cholecalciferol. 1 µg cholecalciferol = 40 IU vitamin D.

^c Under the assumption of minimal sunlight.

^d As α-tocopherol. α-Tocopherol includes *RRR*-α-tocopherol, the only form of α-tocopherol that occurs naturally in foods, and the *2R*-stereoisomeric forms of α-tocopherol (*RRR*-, *RSR*-, *RRS*-, and *RSS*-α-tocopherol) that occur in fortified foods and supplements. It does not include the *2S*-stereoisomeric forms of α-tocopherol (*SRR*-, *SSR*-, *SRS*-, and *SSS*-α-tocopherol), also found in fortified foods and supplements.

^e As niacin equivalents (NE). 1 mg of niacin = 60 mg of tryptophan; 0–6 months = preformed niacin (not NE).

^f As dietary folate equivalents (DFE). 1 DFE = 1 µg food folate = 0.6 µg of folic acid from fortified food or as a supplement consumed with food = 0.5 µg of a supplement taken on an empty stomach.

^g Although AIs have been set for choline, there are few data to assess whether a dietary supply of choline is needed at all stages of the life cycle, and it may be that the choline requirement can be met by endogenous synthesis at some of these stages.

^h Because 10 to 30 percent of older people may malabsorb food-bound B₁₂, it is advisable for those older than 50 years to meet their RDA mainly by consuming foods fortified with B₁₂ or a supplement containing B₁₂.

ⁱ In view of evidence linking folate intake with neural tube defects in the fetus, it is recommended that all women capable of becoming pregnant consume 400 µg from supplements or fortified foods in addition to intake of food folate from a varied diet.

Dietary Reference Intakes (DRIs): Recommended Dietary Allowances and Adequate Intakes, Elements

Food and Nutrition Board, Institute of Medicine, National Academies

Life Stage Group	Calcium (mg/d)	Chromium (µg/d)	Copper (µg/d)	Fluoride (mg/d)	Iodine (µg/d)	Iron (mg/d)	Magnesium (mg/d)	Manganese (mg/d)	Molybdenum (µg/d)	Phosphorus (mg/d)	Selenium (µg/d)	Zinc (mg/d)	Potassium (g/d)	Sodium (g/d)	Chloride (g/d)
Infants															
0 to 6 mo	200*	0.2*	200*	0.01*	110*	0.27*	30*	0.003*	2*	100*	15*	2*	0.4*	0.12*	0.18*
6 to 12 mo	260*	5.5*	220*	0.5*	130*	11	75*	0.6*	3*	275*	20*	3	0.7*	0.37*	0.57*
Children															
1–3 y	700	11*	340	0.7*	90	7	80	1.2*	17	460	20	3	3.0*	1.0*	1.5*
4–8 y	1,000	15*	440	1*	90	10	130	1.5*	22	500	30	5	3.8*	1.2*	1.9*
Males															
9–13 y	1,300	25*	700	2*	120	8	240	1.9*	34	1,250	40	8	4.5*	1.5*	2.3*
14–18 y	1,300	35*	890	3*	150	11	410	2.2*	43	1,250	55	11	4.7*	1.5*	2.3*
19–30 y	1,000	35*	900	4*	150	8	400	2.3*	45	700	55	11	4.7*	1.5*	2.3*
31–50 y	1,000	35*	900	4*	150	8	420	2.3*	45	700	55	11	4.7*	1.5*	2.3*
51–70 y	1,000	30*	900	4*	150	8	420	2.3*	45	700	55	11	4.7*	1.3*	2.0*
> 70 y	1,200	30*	900	4*	150	8	420	2.3*	45	700	55	11	4.7*	1.2*	1.8*
Females															
9–13 y	1,300	21*	700	2*	120	8	240	1.6*	34	1,250	40	8	4.5*	1.5*	2.3*
14–18 y	1,300	24*	890	3*	150	15	360	1.6*	43	1,250	55	9	4.7*	1.5*	2.3*
19–30 y	1,000	25*	900	3*	150	18	310	1.8*	45	700	55	8	4.7*	1.5*	2.3*
31–50 y	1,000	25*	900	3*	150	18	320	1.8*	45	700	55	8	4.7*	1.5*	2.3*
51–70 y	1,200	20*	900	3*	150	8	320	1.8*	45	700	55	8	4.7*	1.3*	2.0*
> 70 y	1,200	20*	900	3*	150	8	320	1.8*	45	700	55	8	4.7*	1.2*	1.8*
Pregnancy															
14–18 y	1,300	29*	1,000	3*	220	27	400	2.0*	50	1,250	60	12	4.7*	1.5*	2.3*
19–30 y	1,000	30*	1,000	3*	220	27	350	2.0*	50	700	60	11	4.7*	1.5*	2.3*
31–50 y	1,000	30*	1,000	3*	220	27	360	2.0*	50	700	60	11	4.7*	1.5*	2.3*
Lactation															
14–18 y	1,300	44*	1,300	3*	290	10	360	2.6*	50	1,250	70	13	5.1*	1.5*	2.3*
19–30 y	1,000	45*	1,300	3*	290	9	310	2.6*	50	700	70	12	5.1*	1.5*	2.3*
31–50 y	1,000	45*	1,300	3*	290	9	320	2.6*	50	700	70	12	5.1*	1.5*	2.3*

NOTE: This table (taken from the DRI reports, see www.nap.edu) presents Recommended Dietary Allowances (RDAs) in **bold type** and Adequate Intakes (AIs) in ordinary type followed by an asterisk (*). An RDA is the average daily dietary intake level; sufficient to meet the nutrient requirements of nearly all (97-98 percent) healthy individuals in a group. It is calculated from an Estimated Average Requirement (EAR). If sufficient scientific evidence is not available to establish an EAR, and thus calculate an RDA, an AI is usually developed. For healthy breastfed infants, an AI is the mean intake. The AI for other life stage and gender groups is believed to cover the needs of all healthy individuals in the groups, but lack of data or uncertainty in the data prevent being able to specify with confidence the percentage of individuals covered by this intake.

SOURCES: *Dietary Reference Intakes for Calcium, Phosphorus, Magnesium, Vitamin D, and Fluoride* (1997); *Dietary Reference Intakes for Thiamin, Riboflavin, Niacin, Vitamin B₆, Folate, Vitamin B₁₂, Pantothenic Acid, Biotin, and Choline* (1998); *Dietary Reference Intakes for Vitamin C, Vitamin E, Selenium, and Carotenoids* (2000); and *Dietary Reference Intakes for Vitamin A, Vitamin K, Arsenic, Boron, Chromium, Copper, Iodine, Iron, Manganese, Molybdenum, Nickel, Silicon, Vanadium, and Zinc* (2001); *Dietary Reference Intakes for Water, Potassium, Sodium, Chloride, and Sulfate* (2005); and *Dietary Reference Intakes for Calcium and Vitamin D* (2011). These reports may be accessed via www.nap.edu.

Dietary Reference Intakes (DRIs): Recommended Dietary Allowances and Adequate Intakes, Total Water and Macronutrients

Food and Nutrition Board, Institute of Medicine, National Academies

Life Stage Group	Total Water ^a (L/d)	Carbohydrate (g/d)	Total** Fiber (g/d)	Fat (g/d)	Linoleic Acid (g/d)	α-Linolenic Acid (g/d)	Protein ^b (g/d)
Infants							
0 to 6 mo	0.7*	60*	ND	31*	4.4*	0.5*	9.1*
6 to 12 mo	0.8*	95*	ND	30*	4.6*	0.5*	11.0
Children							
1–3 y	1.3*	130	19*	ND ^c	7*	0.7*	13
4–8 y	1.7*	130	25*	ND	10*	0.9*	19
Males							
9–13 y	2.4*	130	31*	ND	12*	1.2*	34
14–18 y	3.3*	130	38*	ND	16*	1.6*	52
19–30 y	3.7*	130	38*	ND	17*	1.6*	56
31–50 y	3.7*	130	38*	ND	17*	1.6*	56
51–70 y	3.7*	130	30*	ND	14*	1.6*	56
> 70 y	3.7*	130	30*	ND	14*	1.6*	56
Females							
9–13 y	2.1*	130	26*	ND	10*	1.0*	34
14–18 y	2.3*	130	26*	ND	11*	1.1*	46
19–30 y	2.7*	130	25*	ND	12*	1.1*	46
31–50 y	2.7*	130	25*	ND	12*	1.1*	46
51–70 y	2.7*	130	21*	ND	11*	1.1*	46
> 70 y	2.7*	130	21*	ND	11*	1.1*	46
Pregnancy							
14–18 y	3.0*	175	28*	ND	13*	1.4*	71
19–30 y	3.0*	175	28*	ND	13*	1.4*	71
31–50 y	3.0*	175	28*	ND	13*	1.4*	71
Lactation							
14–18	3.8*	210	29*	ND	13*	1.3*	71
19–30 y	3.8*	210	29*	ND	13*	1.3*	71
31–50 y	3.8*	210	29*	ND	13*	1.3*	71

NOTE: This table (take from the DRI reports, see www.nap.edu) presents Recommended Dietary Allowances (RDA) in **bold type** and Adequate Intakes (AI) in ordinary type followed by an asterisk (*). An RDA is the average daily dietary intake level; sufficient to meet the nutrient requirements of nearly all (97-98 percent) healthy individuals in a group. It is calculated from an Estimated Average Requirement (EAR). If sufficient scientific evidence is not available to establish an EAR, and thus calculate an RDA, an AI is usually developed. For healthy breastfed infants, an AI is the mean intake. The AI for other life stage and gender groups is believed to cover the needs of all healthy individuals in the groups, but lack of data or uncertainty in the data prevent being able to specify with confidence the percentage of individuals covered by this intake.

^a Total water includes all water contained in food, beverages, and drinking water.

^b Based on g protein per kg of body weight for the reference body weight, e.g., for adults 0.8 g/kg body weight for the reference body weight.

^cNot determined.

SOURCE: *Dietary Reference Intakes for Energy, Carbohydrate, Fiber, Fat, Fatty Acids, Cholesterol, Protein, and Amino Acids* (2002/2005) and *Dietary Reference Intakes for Water, Potassium, Sodium, Chloride, and Sulfate* (2005). The report may be accessed via www.nap.edu.

****BYU-I NOTE:** The adequate intake (AI) for Total Fiber is based on data which showed that 14 grams per 1000 calories (14g/1000 calories) reduced the risk of coronary heart disease. Age and gender-specific AIs were set by taking 14g/1000 calories and multiplying it by the median energy intake for each group. For example, you will find the Total Fiber AI on the DRI tables (above) for adult men and adult women ages 19-50 is 38 g/day and 25 g/day, respectively. However, some diet analysis programs, including **NutritionCal Plus**, recommend individual Total Fiber intake based on the 14g/1000 calories and multiply it by the individual's recommended calorie intake. Either AI recommendation, the age and gender-specific recommendation or the calorie-based recommendation, can be used and is helpful to ensure adequate consumption of fiber.

Dietary Reference Intakes (DRIs): Acceptable Macronutrient Distribution Ranges

Food and Nutrition Board, Institute of Medicine, National Academies

Macronutrient	Range (percent of energy)		
	Children, 1–3 y	Children, 4–18 y	Adults
Fat	30–40	25–35	20–35
<i>n</i> -6 polyunsaturated fatty acids ^a (linoleic acid)	5–10	5–10	5–10
<i>n</i> -3 polyunsaturated fatty acids ^a (α -linolenic acid)	0.6–1.2	0.6–1.2	0.6–1.2
Carbohydrate	45–65	45–65	45–65
Protein	5–20	10–30	10–35

^a Approximately 10 percent of the total can come from longer-chain *n*-3 or *n*-6 fatty acids.

SOURCE: *Dietary Reference Intakes for Energy, Carbohydrate, Fiber, Fat, Fatty Acids, Cholesterol, Protein, and Amino Acids* (2002/2005). The report may be accessed via www.nap.edu.

Dietary Reference Intakes (DRIs): Acceptable Macronutrient Distribution Ranges

Food and Nutrition Board, Institute of Medicine, National Academies

Macronutrient	Recommendation
Dietary cholesterol	As low as possible while consuming a nutritionally adequate diet
Trans fatty Acids	As low as possible while consuming a nutritionally adequate diet
Saturated fatty acids	As low as possible while consuming a nutritionally adequate diet
Added sugars ^a	Limit to no more than 25 % of total energy

^aNot a recommended intake. A daily intake of added sugars that individuals should aim for to achieve a healthful diet was not set.

SOURCE: *Dietary Reference Intakes for Energy, Carbohydrate, Fiber, Fat, Fatty Acids, Cholesterol, Protein, and Amino Acids* (2002/2005). The report may be accessed via www.nap.edu.

Dietary Reference Intakes (DRIs): Tolerable Upper Intake Levels, Vitamins

Food and Nutrition Board, Institute of Medicine, National Academies

Life Stage Group	Vitamin A (µg/d) ^a	Vitamin C (mg/d)	Vitamin D (µg/d)	Vitamin E (mg/d) ^{b,c}	Vitamin K	Thia-min	Ribo-flavin	Niacin (mg/d) ^f	Vitamin B ₆ (mg/d)	Folate (µg/d) ^c	Vitamin B ₁₂	Panto-thenic Acid	Bio-tin	Cho-line (g/d)	Carote-noids ^d
Infants															
0 to 6 mo	600	ND ^e	25	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
6 to 12 mo	600	ND	38	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
Children															
1–3 y	600	400	63	200	ND	ND	ND	10	30	300	ND	ND	ND	1.0	ND
4–8 y	900	650	75	300	ND	ND	ND	15	40	400	ND	ND	ND	1.0	ND
Males															
9–13 y	1,700	1,200	100	600	ND	ND	ND	20	60	600	ND	ND	ND	2.0	ND
14–18 y	2,800	1,800	100	800	ND	ND	ND	30	80	800	ND	ND	ND	3.0	ND
19–30 y	3,000	2,000	100	1,000	ND	ND	ND	35	100	1,000	ND	ND	ND	3.5	ND
31–50 y	3,000	2,000	100	1,000	ND	ND	ND	35	100	1,000	ND	ND	ND	3.5	ND
51–70 y	3,000	2,000	100	1,000	ND	ND	ND	35	100	1,000	ND	ND	ND	3.5	ND
> 70 y	3,000	2,000	100	1,000	ND	ND	ND	35	100	1,000	ND	ND	ND	3.5	ND
Females															
9–13 y	1,700	1,200	100	600	ND	ND	ND	20	60	600	ND	ND	ND	2.0	ND
14–18 y	2,800	1,800	100	800	ND	ND	ND	30	80	800	ND	ND	ND	3.0	ND
19–30 y	3,000	2,000	100	1,000	ND	ND	ND	35	100	1,000	ND	ND	ND	3.5	ND
31–50 y	3,000	2,000	100	1,000	ND	ND	ND	35	100	1,000	ND	ND	ND	3.5	ND
51–70 y	3,000	2,000	100	1,000	ND	ND	ND	35	100	1,000	ND	ND	ND	3.5	ND
> 70 y	3,000	2,000	100	1,000	ND	ND	ND	35	100	1,000	ND	ND	ND	3.5	ND
Pregnancy															
14–18 y	2,800	1,800	100	800	ND	ND	ND	30	80	800	ND	ND	ND	3.0	ND
19–30 y	3,000	2,000	100	1,000	ND	ND	ND	35	100	1,000	ND	ND	ND	3.5	ND
31–50 y	3,000	2,000	100	1,000	ND	ND	ND	35	100	1,000	ND	ND	ND	3.5	ND
Lactation															
14–18 y	2,800	1,800	100	800	ND	ND	ND	30	80	800	ND	ND	ND	3.0	ND
19–30 y	3,000	2,000	100	1,000	ND	ND	ND	35	100	1,000	ND	ND	ND	3.5	ND
31–50 y	3,000	2,000	100	1,000	ND	ND	ND	35	100	1,000	ND	ND	ND	3.5	ND

NOTE: A Tolerable Upper Intake Level (UL) is the highest level of daily nutrient intake that is likely to pose no risk of adverse health effects to almost all individuals in the general population. Unless otherwise specified, the UL represents total intake from food, water, and supplements. Due to a lack of suitable data, ULs could not be established for vitamin K, thiamin, riboflavin, vitamin B₁₂, pantothenic acid, biotin, and carotenoids. In the absence of a UL, extra caution may be warranted in consuming levels above recommended intakes. Members of the general population should be advised not to routinely exceed the UL. The UL is not meant to apply to individuals who are treated with the nutrient under medical supervision or to individuals with predisposing conditions that modify their sensitivity to the nutrient.

^aAs preformed vitamin A only.

^bAs α-tocopherol; applies to any form of supplemental α-tocopherol.

^cThe ULs for vitamin E, niacin, and folate apply to synthetic forms obtained from supplements, fortified foods, or a combination of the two.

^dβ-Carotene supplements are advised only to serve as a provitamin A source for individuals at risk of vitamin A deficiency.

^eND = Not determinable due to lack of data of adverse effects in this age group and concern with regard to lack of ability to handle excess amounts. Source of intake should be from food only to prevent high levels of intake.

SOURCES: *Dietary Reference Intakes for Calcium, Phosphorous, Magnesium, Vitamin D, and Fluoride* (1997); *Dietary Reference Intakes for Thiamin, Riboflavin, Niacin, Vitamin B₆, Folate, Vitamin B₁₂, Pantothenic Acid, Biotin, and Choline* (1998); *Dietary Reference Intakes for Vitamin C, Vitamine E, Selenium, and Carotenoids* (2000); *Dietary Reference Intakes for Vitamin A, Vitamin K, Arsenic, Boron, Chromium, Copper, Iodine, Iron, Manganese, Molybdenum, Nickel, Silicon, Vanadium, and Zinc* (2001); and *Dietary Reference Intakes for Calcium and Vitamin D* (2011). These reports may be accessed via www.nap.edu.

Dietary Reference Intakes (DRIs): Tolerable Upper Intake Levels, Elements

Food and Nutrition Board, Institute of Medicine, National Academies

Life Stage Group	Arsenic ^a	Boron (mg/d)	Calcium (mg/d)	Chromium	Copper (µg/d)	Fluoride (mg/d)	Iodine (µg/d)	Iron (mg/d)	Magnesium (mg/d) ^b	Manganese (mg/d)	Molybdenum (µg/d)	Nickel (mg/d)	Phosphorus (g/d)	Selenium (µg/d)	Silicon ^c	Vanadium (mg/d) ^d	Zinc (mg/d)	Sodium (g/d)	Chloride (g/d)
Infants																			
0 to 6 mo	ND ^e	ND	1,000	ND	ND	0.7	ND	40	ND	ND	ND	ND	ND	45	ND	ND	4	ND	ND
6 to 12 mo	ND	ND	1,500	ND	ND	0.9	ND	40	ND	ND	ND	ND	ND	60	ND	ND	5	ND	ND
Children																			
1–3 y	ND	3	2,500	ND	1,000	1.3	200	40	65	2	300	0.2	3	90	ND	ND	7	1.5	2.3
4–8 y	ND	6	2,500	ND	3,000	2.2	300	40	110	3	600	0.3	3	150	ND	ND	12	1.9	2.9
Males																			
9–13 y	ND	11	3,000	ND	5,000	10	600	40	350	6	1,100	0.6	4	280	ND	ND	23	2.2	3.4
14–18 y	ND	17	3,000	ND	8,000	10	900	45	350	9	1,700	1.0	4	400	ND	ND	34	2.3	3.6
19–30 y	ND	20	2,500	ND	10,000	10	1,100	45	350	11	2,000	1.0	4	400	ND	1.8	40	2.3	3.6
31–50 y	ND	20	2,500	ND	10,000	10	1,100	45	350	11	2,000	1.0	4	400	ND	1.8	40	2.3	3.6
51–70 y	ND	20	2,000	ND	10,000	10	1,100	45	350	11	2,000	1.0	4	400	ND	1.8	40	2.3	3.6
> 70 y	ND	20	2,000	ND	10,000	10	1,100	45	350	11	2,000	1.0	3	400	ND	1.8	40	2.3	3.6
Females																			
9–13 y	ND	11	3,000	ND	5,000	10	600	40	350	6	1,100	0.6	4	280	ND	ND	23	2.2	3.4
14–18 y	ND	17	3,000	ND	8,000	10	900	45	350	9	1,700	1.0	4	400	ND	ND	34	2.3	3.6
19–30 y	ND	20	2,500	ND	10,000	10	1,100	45	350	11	2,000	1.0	4	400	ND	1.8	40	2.3	3.6
31–50 y	ND	20	2,500	ND	10,000	10	1,100	45	350	11	2,000	1.0	4	400	ND	1.8	40	2.3	3.6
51–70 y	ND	20	2,000	ND	10,000	10	1,100	45	350	11	2,000	1.0	4	400	ND	1.8	40	2.3	3.6
> 70 y	ND	20	2,000	ND	10,000	10	1,100	45	350	11	2,000	1.0	3	400	ND	1.8	40	2.3	3.6
Pregnancy																			
14–18 y	ND	17	3,000	ND	8,000	10	900	45	350	9	1,700	1.0	3.5	400	ND	ND	34	2.3	3.6
19–30 y	ND	20	2,500	ND	10,000	10	1,100	45	350	11	2,000	1.0	3.5	400	ND	ND	40	2.3	3.6
61–50 y	ND	20	2,500	ND	10,000	10	1,100	45	350	11	2,000	1.0	3.5	400	ND	ND	40	2.3	3.6
Lactation																			
14–18 y	ND	17	3,000	ND	8,000	10	900	45	350	9	1,700	1.0	4	400	ND	ND	34	2.3	3.6
19–30 y	ND	20	2,500	ND	10,000	10	1,100	45	350	11	2,000	1.0	4	400	ND	ND	40	2.3	3.6
31–50 y	ND	20	2,500	ND	10,000	10	1,100	45	350	11	2,000	1.0	4	400	ND	ND	40	2.3	3.6

NOTE: A Tolerable Upper Intake Level (UL) is the highest level of daily nutrient intake that is likely to pose no risk of adverse health effects to almost all individuals in the general population. Unless otherwise specified, the UL represents total intake from food, water, and supplements. Due to a lack of suitable data, ULs could not be established for vitamin K, thiamin, riboflavin, vitamin B₁₂, pantothenic acid, biotin, and carotenoids. In the absence of a UL, extra caution may be warranted in consuming levels above recommended intakes. Members of the general population should be advised not to routinely exceed the UL. The UL is not meant to apply to individuals who are treated with the nutrient under medical supervision or to individuals with predisposing conditions that modify their sensitivity to the nutrient.

^aAlthough the UL was not determined for arsenic, there is no justification for adding arsenic to food or supplements.

^bThe ULs for magnesium represent intake from a pharmacological agent only and do not include intake from food and water.

^cAlthough silicon has not been shown to cause adverse effects in humans, there is no justification for adding silicon to supplements.

^dAlthough vanadium in food has not been shown to cause adverse effects in humans, there is no justification for adding vanadium to food and vanadium supplements should be used with caution. The UL is based on adverse effects in laboratory animals and this data could be used to set a UL for adults but not children and adolescents.

^eND = Not determinable due to lack of data of adverse effects in this age group and concern with regard to lack of ability to handle excess amounts. Source of intake should be from food only to prevent high levels of intake.

SOURCES: *Dietary Reference Intakes for Calcium, Phosphorous, Magnesium, Vitamin D, and Fluoride* (1997); *Dietary Reference Intakes for Thiamin, Riboflavin, Niacin, Vitamin B₆, Folate, Vitamin B₁₂, Pantothenic Acid, Biotin, and Choline* (1998); *Dietary Reference Intakes for Vitamin C, Vitamin E, Selenium, and Carotenoids* (2000); *Dietary Reference Intakes for Vitamin A, Vitamin K, Arsenic, Boron, Chromium, Copper, Iodine, Iron, Manganese, Molybdenum, Nickel, Silicon, Vanadium, and Zinc* (2001); *Dietary Reference Intakes for Water, Potassium, Sodium, Chloride, and Sulfate* (2005); and *Dietary Reference Intakes for Calcium and Vitamin D* (2011). These reports may be accessed via www.nap.edu.